

Gravity and Light: Combining Gravitational Wave and Electromagnetic Observations in the 2020s

Thematic areas: 3. Stars and Stellar Evolution 4. Formation and Evolution of Compact Objects 7. Cosmology and Fundamental Physics 8. Multi-Messenger Astronomy and Astrophysics

Principal Author:

Ryan J. Foley
UC Santa Cruz
foley@ucsc.edu

Co-authors:

K. D. Alexander¹, I. Andreoni², I. Arcavi³, K. Auchettl⁴, J. Barnes⁵, G. Baym⁶, E. C. Bellm⁷, A. M. Beloborodov⁵, N. Blagorodnova⁸, J. P. Blakeslee⁹, P. R. Brady¹⁰, M. Branchesi^{11,12}, J. S. Brown¹³, N. Butler¹⁴, M. Cantiello¹⁵, R. Chornock¹⁶, D. O. Cook^{2,17}, J. Cooke¹⁸, D. L. Coppejans¹, A. Corsi¹⁹, S. M. Couch²⁰, M. W. Coughlin², D. A. Coulter¹³, P. S. Cowperthwaite²¹, T. Dietrich²², G. Dimitriadis¹³, M. R. Drout²³, J. H. Elias^{24,25}, B. Farr²⁶, R. Fernandez²⁷, A. V. Filippenko²⁸, W. Fong¹, T. Fragos²⁹, D. A. Frail³⁰, W. L. Freedman³¹, C. L. Fryer³², V. Z. Golkhou⁷, D. Hiramatsu^{33,34}, J. Hjorth^{4,35}, A. Hoshino³⁶, G. Hosseinzadeh³⁷, K. Hotokezaka³⁸, D. A. Howell^{33,34}, T. Hung¹³, D. O. Jones¹³, V. Kalogera¹, D. Kasen²⁸, W. E. Kerzendorf³⁹, C. D. Kilpatrick¹³, R. P. Kirshner³⁷, K. Krisciunas⁴⁰, J. M. Lattimer⁴¹, D. Lazzati⁴², A. J. Levan⁸, A. I. MacFadyen³⁹, K. Maeda⁴³, I. Mandel⁴⁴, K. S. Mandel⁴⁵, B. Margalit²⁸, R. Margutti¹, J. McIver², B. D. Metzger⁵, M. Modjaz³⁹, K. Mooley^{30,2}, T. Moriya⁴⁶, A. Murguia-Berthier¹³, G. Narayan⁴⁷, M. Nicholl⁴⁸, S. Nissanke⁴⁹, K. Nomoto⁵⁰, J. M. O’Meara⁵¹, R. O’Shaughnessy⁵², E. O’Connor⁵³, A. Palmese⁵⁴, Y.-C. Pan⁴⁶, C. Pankow¹, K. Paterson¹, D. A. Perley⁵⁵, R. Perna⁴¹, A. L. Piro²¹, T. A. Pritchard³⁹, E. Quataert²⁸, D. Radice³⁸, E. Ramirez-Ruiz¹³, S. Reddy⁷, A. Rest⁴⁷, A. G. Riess⁵⁶, C. L. Rodriguez⁵⁷, C. Rojas-Bravo¹³, E. M. Rossi⁵⁸, S. Rosswog⁵³, M. Ruiz⁶, S. L. Shapiro⁶, D. H. Shoemaker⁵⁷, M. R. Siebert¹³, D. M. Siegel⁵, K. Siellez¹³, N. Smith⁵⁹, M. Soares-Santos⁶⁰, N. B. Suntzeff⁴⁰, R. Surman⁶¹, M. Tanaka⁶², N. R. Tanvir⁶³, G. Terreran¹, S. Valenti⁶⁴, V. A. Villar³⁷, L. Wang⁴⁰, S. A. Webb⁶⁵, J. C. Wheeler⁶⁶, P. K. G. Williams^{67,68}, S. Woosley¹³, M. Zaldarriaga⁶⁹, M. Zevin¹

¹Northwestern, ²Caltech, ³Tel Aviv, ⁴DARK, ⁵Columbia, ⁶UIUC, ⁷UW, ⁸Raboud, ⁹Gemini, ¹⁰UW-Milwaukee, ¹¹GSII, ¹²INFN, ¹³UC Santa Cruz, ¹⁴ASU, ¹⁵CCA, ¹⁶Ohio, ¹⁷IPAC, ¹⁸Swinburne, ¹⁹TTU, ²⁰MSU, ²¹Carnegie, ²²NIKHEF, ²³Toronto, ²⁴SOAR, ²⁵NOAO, ²⁶Oregon, ²⁷Alberta, ²⁸UC Berkeley, ²⁹Geneva, ³⁰NRAO, ³¹Chicago, ³²LANL, ³³Las Cumbres, ³⁴UC Santa Barbara, ³⁵NBI, ³⁶HUJI, ³⁷Harvard, ³⁸Princeton, ³⁹NYU, ⁴⁰TAMU, ⁴¹Stony Brook, ⁴²Oregon State, ⁴³Kyoto, ⁴⁴Monash, ⁴⁵Cambridge, ⁴⁶NAOJ, ⁴⁷STScI, ⁴⁸Edinburgh, ⁴⁹Amsterdam, ⁵⁰IPMU, ⁵¹Keck, ⁵²RIT, ⁵³Stockholm, ⁵⁴FNAL, ⁵⁵LJMU, ⁵⁶JHU, ⁵⁷MIT, ⁵⁸Leiden, ⁵⁹Arizona, ⁶⁰Brandeis, ⁶¹Notre Dame, ⁶²Tohoku, ⁶³Leicester, ⁶⁴UC Davis, ⁶⁵SUT, ⁶⁶UT Austin, ⁶⁷CfA, ⁶⁸AAS, ⁶⁹IAS

Abstract:

As of today, we have directly detected exactly one source in both gravitational waves (GWs) and electromagnetic (EM) radiation, the binary neutron star merger GW170817, its associated gamma-ray burst GRB170817A, and the subsequent kilonova SSS17a/AT 2017gfo. Within ten years, we will detect hundreds of events, including new classes of events such as neutron-star–black-hole mergers, core-collapse supernovae, and almost certainly something completely unexpected. As we build this sample, we will explore exotic astrophysical topics ranging from nucleosynthesis, stellar evolution, general relativity, high-energy astrophysics, nuclear matter, to cosmology. The discovery potential is extraordinary, and investments in this area will yield major scientific breakthroughs. Here we outline some of the most exciting scientific questions that can be answered by combining GW and EM observations.

1 The Dawn of Multi-Messenger Astronomy

On August 17, 2017, the LIGO-Virgo collaboration (LVC) detected, for the first time, a gravitational wave (GW) signal (GW170817) from two inspiralling neutron stars [NSs; 6]. At the same time an electromagnetic (EM) counterparts, a short gamma-ray burst (SGRB; GRB170817A) and a kilonova (called SSS17a or AT 2017gfo), was detected at all wavelengths [5, 17, 30, 75]. With a single event, we were able to address fundamental questions in general relativity [56], nuclear matter [6], cosmology [4, 19, 24], nucleosynthesis [36], and astrophysics [e.g., 18, 22, 36, 38, 40, 58, 72].

In April 2019, LVC will start its third observing run, O3, at increased sensitivity. O3 will last about a year (roughly through March 2020). LVC expects to detect between 1 and 50 binary NS (BNS) mergers during O3 with a median estimate of about 10 events [2]. There is also the possibility of discovering the first neutron-star–black-hole (NSBH) merger, which should produce an EM counterpart similar to (but likely fainter than) those of BNS mergers. Toward the end of O3, KAGRA may join the network, and with it, we will have our first 4-detector events. In 2021, LVC (with KAGRA) is expected to start another observing run at design sensitivity [3]. LIGO India should join the network in 2025. Over the next decade, we will move from having a single event seen in both gravitational waves and light to a sample of hundreds.

As this new field quickly evolves, we are poised to use gravitational waves and light in concert to answer long-standing questions in nucleosynthesis, stellar evolution, nuclear matter, and cosmology. **We are confident that the combination of gravity and light will allow us to make major strides in several vital areas, including generating surprises.**

2 What are the Origins of the Heaviest Elements?

The light curves and spectra of SSS17a indicate that up to $0.05 M_{\odot}$ of r -process material was created in and ejected from the merger. Combined with the BNS merger rate, BNS mergers could produce all of the r -process material in our Galaxy *if* SSS17a is typical [18, 40, 66, 67, 72, 73]. However, the total yield per kilonova depends on the mass ratio of the binary and the total remnant mass. Studies of emission from SGRBs indicate that GW170817 produced more r -process material than the typical SGRB [10, 29].

Even in the case of SSS17a, the exact amount of r -process material, and particularly “heavy” or “3rd-peak” material, is still quite uncertain [78]. The optical and infrared radiation in kilonovae can arise from a variety of ejecta components and even radioactive species in the merger, tidal tails, and accretion disk winds [36, 50], and precise mass estimates require a full understanding of the emission mechanisms and careful accounting. For SSS17a it is still debated whether the color evolution was caused by (1) decay of r -process nuclei in ejecta with different compositions [39, 76]; (2) decay of material from a single origin [72, 77]; or (3) energy sources such as shock cooling [59], remnant winds [79], or central engine activity [41]. Ultraviolet through infrared light curves of a statistical sample of kilonovae will allow us to distinguish between models and address key theoretical questions.

An intriguing path toward measuring the amount of heavy r -process material is through careful examination of the kilonova light curve at late times. At early times, many radioactive species contribute to the luminosity and the combination of all decays results in a power-

law-dependent heating rate. At later times, as the majority of species decay away, the kilonova light curve is powered by at most a handful of long-lived radioisotopes. Theorists have identified ^{254}Cf , a heavy r -process element with a half-life of 60.5 days [78, 80], as being a possible dominant energy source at late times. Measuring the kilonova luminosity two months after the merger will reveal the amount of ^{254}Cf generated in the merger [37]. Combined with the robust solar abundance pattern and the early-time data, we can then accurately determine the relative abundance of light-to-heavy r -process material created in BNS mergers.

Perhaps the most compelling path toward further understanding element creation is through spectral modeling [36]. Currently, atomic line lists are insufficient for directly determining the abundance of individual elements in kilonova spectra [35]. Improvements in this area will require significant laboratory work, but it would be extremely beneficial to this science area. Better spectral modeling will also strengthen our constraints on the dynamics of the ejecta, especially once we obtain a nebular spectrum with JWST or an ELT.

3 What are the Details of the Merger and Explosion?

Numerical relativity can precisely reproduce the waveform for the inspiral before a BNS merger [6]. However, LVC was unable to detect the actual merger for GW170817 since its frequency increased to a point where its detectors lose sensitivity. As a result, we must rely on numerical simulations to understand the actual merger process [27, 42, 57, 65, 68]. These simulations generically show that the NSs should be (partially) tidally disrupted as they merge. Depending on the individual NS masses and the mass ratio of the system, there is a broad range of outcomes. The ejecta masses, ejecta velocities, number of tidal tails, and duration/size of an accretion disk all depend on the details of the progenitor system.

The merger product could promptly collapse to a BH [11] or could be an accreting hypermassive NS (HMNS) that exists for several milliseconds before collapsing to a black hole [71]. If the latter, a significant neutrino flux will exist during this time, which can irradiate the disk wind, changing the electron fraction and thus the final composition [22, 23, 34, 40, 49]. The size of the disk and the lifetime of the HMNS both depend on the equation-of-state (EoS) for nuclear matter [32]. Since the composition of the ejecta affects the EM observables, one can hope to invert the problem to understand the immediate aftermath of the merger [12].

However, additional physics may complicate the picture. In particular, we expect a relativistic jet to be launched from at least some BNS mergers. If the jet escapes, it should heat a cocoon of material surrounding the jet that can produce additional early-time emission. Shock cooling, a long-lived engine, and additional disk-wind emission could all contribute to the early-time flux. Detailed modeling of these processes and comparison to the non-thermal emission at γ -ray, X-ray, UV, and radio wavelengths, especially observations within the first day after merger as well as months later, should differentiate between scenarios [7, 9, 28, 44, 52].

The combination of GW and EM data will be particularly useful for compact binary coalescences. The GW signal precisely determines the merger time, removing a key variable from modeling the EM data. For these events the GW data precisely measures the binary chirp mass, but the mass ratio is not as well constrained. GW data also set upper limits to the tidal deformations and thus the radii of NSs. EM data complement these analyses,

as these data in combination with numerical relativity simulations place lower limits on the tidal deformation and more stringent constraints on the mass ratio. Independently, the EM signal depends on the maximum mass of NSs [48], a unique look at the EOS at the center of the NS. In this way, combined GW and EM data allow for a better understanding of the outcomes and physical processes of compact binary coalescence mergers [e.g., 16, 21, 63]

4 What Systems Create Compact-Object Mergers?

GW observations of binary BHs have revealed the population of massive stellar-mass BHs [20–50 M_{\odot} ; 74], raising questions about which stars produce the BHs, how such systems form, and what the minimum and maximum masses are for stellar-mass black holes. Similar questions exist for systems that include at least one NS, but with even fewer data. While we have observed a limited number of BNS systems in the Milky Way and have some constraints from short gamma-ray bursts (SGRBs), both populations must be biased subsamples of the full population of compact-object binaries.

As we build up statistical samples of compact-object mergers, we will be able to examine their demographics, revealing details of the underlying population. As an example, the SGRB host-galaxy population is skewed toward star-forming galaxies, but a significant fraction still resides in passive galaxies [e.g., 26, 62]. This implies a delay-time distribution (the time from star formation until merger) is peaked toward short delay times, but has a long tail (if gravitational radiation is the slowest step, one expects a rate proportional to t^{-1} , consistent with the observations). Similarly, a sizable fraction of SGRBs are found far from their host galaxies or have no associated host galaxy [25], indicating that these systems were expelled from their birth sites by supernova (SN) kicks. But it is unclear how much the population of SGRBs, especially those with localizations sufficient to determine a host galaxy, overlap with BNS mergers.

We will be able to independently determine the delay-time distribution and offset distribution for a sample originating from GW detections. With detailed analyses of the host environments, we will generate a more explicit link to BNS progenitor systems [13, 31, 45, 54, 55].

We will also be able to directly compare the physical properties determined from the GW signal with both kilonova properties and population statistics. For instance, one might expect systems with large mass ratios to be disrupted if there is a large SN kick and therefore all such systems are located within two half-light radii of their host galaxy.

While answering questions about the underlying population, we will also determine if every BNS merger produces a GRB, if SGRB samples are biased against significantly kicked mergers, and if BNS and NSBH mergers produce similar gamma-ray observables.

5 How Fast is the Universe Expanding?

New, independent measurements of H_0 are especially important since there is currently a $3.8\text{-}\sigma$ disagreement between the local value [64] and that of the inverse-distance ladder [60]. As the H_0 tension is one of the most significant open questions in cosmology and could reveal new physics, including non- Λ dark energy, it is essential to use as many reliable techniques as possible.

Unlike most astrophysical sources, the waveform of GWs directly encodes the luminosity distance [70]. However, once an EM counterpart is found, we can independently measure

a redshift from a host galaxy or the transient itself (which could be especially crucial for events kicked far from their host galaxy). With both the luminosity distance and redshift, we can measure H_0 .

With GW170817, we made the first standard siren measurement of H_0 [4], finding a value consistent with both the local and inverse distance ladder values. The initial analysis only used the position of the source and the redshift of its host galaxy. The constraint on H_0 was limited by the degeneracy between H_0 and the inclination angle of the binary system. The inclination angle measurement was further improved by radio observations of the jet [51], significantly reducing the uncertainty on H_0 [33].

As new events are added to the sample, the precision scales roughly as \sqrt{N} [14, 15, 20, 53]. The exact improvement depends on the details of the system, where some objects with smaller distance uncertainties or very well-constrained inclination angles will be much more valuable than others. Simple predictions indicate that we can make a 3% measurement of H_0 with 40 GW events with counterparts [14], which should be achievable within the next few years.

Finally, once the population has increased, we can use properties of the sample to further improve all measurements. Specifically, BNS mergers should be randomly oriented [although we need to pay careful attention to selection effects; 47]. With a hierarchical model, we can enforce that our sample has an appropriate inclination distribution. This effectively reduces the possible inclination angles for otherwise poorly constrained events.

While systematic uncertainties are currently subdominant, they may become important at the few percent level. Further improving the calibration of the GW strain signal and studies of the merger and kilonova emission will be critical to harness the full statistical sample that we will have in 10 years.

With improved GW facilities, we will be able to detect events sufficiently far to measure $H(z)$ *directly*. Future facilities such as LIGO Voyager and the proposed Cosmic Explorer should be able to detect BNS mergers to redshift $z \approx 0.4$ and $z \approx 7$, respectively¹. Combined with sensitive space-based NIR observatories such as *JWST*, *WFIRST*, *LUVOIR*, and ground-based ELTs, we could directly track the expansion history of the Universe to the age of deceleration, providing an independent probe of dark energy [20].

6 What Will We Learn from Other Sources of Gravitational Radiation?

During the process of core collapse, stellar cores are expected to generate relatively strong gravitational radiation [43]. While not nearly as loud as BNS mergers, the LVC is already sensitive to Milky Way supernovae [with some extreme models being much louder and possibly detectable in other galaxies; 1]. Such an SN in the Milky Way would be especially interesting as we would also expect to detect thousands of associated neutrinos [69].

Although the GW signal from core collapse should be more complicated than binary coalescence, the waveform would encode critical information about the core-collapse process, the proto-NS, and the SN mechanism [46, 61].

As Galactic SNe are relatively rare (there is perhaps a 10% chance of one occurring in the 2020s), it is possible that GW detectors will improve quickly enough that the first GW

¹ https://dcc.ligo.org/public/0150/T1800084/005/gw_horizons.pdf

detection from an SN will be in another galaxy. Future upgrades and new facilities could reliably detect SNe in the Virgo cluster. As there are a handful of core-collapse SNe within 20 Mpc each year, we could have a sample of several by the end of the decade. With such a sample, we would be able to determine if there are multiple explosion mechanisms and if the kinematics of the core is related to the success of an SN, the mass of the star, or the details of the resulting explosion.

While we *expect* to eventually detect SNe with GWs, there could be GW sources not yet theorized. Every other time astronomers have opened a new window to the Universe, there have been new physical phenomena observed. For GWs, the discovery space is particularly large, increasing the anticipation and expectation for something new. EM observations will be critical to localize and characterize any such source.

7 Recommendations

The combination of gravitational wave and electromagnetic observations will be one of the most fruitful scientific endeavors of the next decade. While this scientific field is still in its infancy, the scientific return for a wide range of fields, extending beyond astrophysics, is enormous. To take full advantage of this opportunity, we make the following recommendations.

- Increase the number, sensitivity, frequency range, calibration accuracy, and duty cycle of GW detectors.
- Ensure that all appropriate observatories have responsive target-of-opportunity programs and provide a sufficient fraction of observing time to such programs. Observatories should limit restrictions on the execution of these programs and work toward automation when feasible.
- Incentivize and support building tools to enable data discovery and sharing [8, see also Chang et al. white paper].
- Develop and maintain sensitive IR instruments for photometry and spectroscopy on medium- and large-aperture telescopes on the ground and in space.
- Construct ELTs, which will be critical for obtaining data at late times and for lower-luminosity/distant events, in both hemispheres (see also Chornock et al. white paper).
- Continue to support and operate sensitive γ -ray and X-ray observatories.
- Plan and develop new UV space observatories, with an emphasis on improving the sensitivity over current capabilities.
- Continue to support and improve sensitive radio observatories (e.g., ngVLA).
- Support the application of theory and high-performance computing for the simulations of mergers, SN explosions, and other exciting objects to connect their GW and EM signals.
- Obtain complementary observational (e.g., NICER) and laboratory data and perform additional theoretical modeling of the NS EoS.
- Perform additional laboratory measurements to improve the atomic spectral-line database of heavy elements.
- Enable continuous GW monitoring to ensure detection of the next Galactic SN.

References

- [1] B. P. Abbott, R. Abbott, T. D. Abbott, M. R. Abernathy, F. Acernese, K. Ackley, C. Adams, T. Adams, P. Addesso, R. X. Adhikari, and et al. First targeted search for gravitational-wave bursts from core-collapse supernovae in data of first-generation laser interferometer detectors. , 94(10):102001, Nov. 2016.
- [2] B. P. Abbott, R. Abbott, T. D. Abbott, M. R. Abernathy, F. Acernese, K. Ackley, C. Adams, T. Adams, P. Addesso, R. X. Adhikari, and et al. Prospects for Observing and Localizing Gravitational-Wave Transients with Advanced LIGO and Advanced Virgo. *Living Reviews in Relativity*, 19, Feb. 2016.
- [3] B. P. Abbott, R. Abbott, T. D. Abbott, M. R. Abernathy, F. Acernese, K. Ackley, C. Adams, T. Adams, P. Addesso, R. X. Adhikari, and et al. Prospects for observing and localizing gravitational-wave transients with Advanced LIGO, Advanced Virgo and KAGRA. *Living Reviews in Relativity*, 21:3, Apr. 2018.
- [4] B. P. Abbott, R. Abbott, T. D. Abbott, F. Acernese, K. Ackley, C. Adams, T. Adams, P. Addesso, R. X. Adhikari, V. B. Adya, and et al. A gravitational-wave standard siren measurement of the Hubble constant. *Nature*, 551:85–88, Nov. 2017.
- [5] B. P. Abbott, R. Abbott, T. D. Abbott, F. Acernese, K. Ackley, C. Adams, T. Adams, P. Addesso, R. X. Adhikari, V. B. Adya, and et al. Gravitational Waves and Gamma-Rays from a Binary Neutron Star Merger: GW170817 and GRB 170817A. *ApJL*, 848:L13, Oct. 2017.
- [6] B. P. Abbott, R. Abbott, T. D. Abbott, F. Acernese, K. Ackley, C. Adams, T. Adams, P. Addesso, R. X. Adhikari, V. B. Adya, and et al. GW170817: Observation of Gravitational Waves from a Binary Neutron Star Inspiral. *Physical Review Letters*, 119(16):161101, Oct. 2017.
- [7] K. D. Alexander, R. Margutti, P. K. Blanchard, W. Fong, E. Berger, A. Hajela, T. Eftekhari, R. Chornock, P. S. Cowperthwaite, D. Giannios, C. Guidorzi, A. Kathirgamaraju, A. MacFadyen, B. D. Metzger, M. Nicholl, L. Sironi, V. A. Villar, P. K. G. Williams, X. Xie, and J. Zrake. A Decline in the X-Ray through Radio Emission from GW170817 Continues to Support an Off-axis Structured Jet. *ApJL*, 863:L18, Aug. 2018.
- [8] G. Allen, I. Andreoni, E. Bachelet, G. B. Berriman, F. B. Bianco, R. Biswas, M. Carasco Kind, K. Chard, M. Cho, P. S. Cowperthwaite, Z. B. Etienne, D. George, T. Gibbs, M. Graham, W. Gropp, A. Gupta, R. Haas, E. A. Huerta, E. Jennings, D. S. Katz, A. Khan, V. Kindratenko, W. T. C. Kramer, X. Liu, A. Mahabal, K. McHenry, J. M. Miller, M. S. Neubauer, S. Oberlin, A. R. Olivas, Jr, S. Rosofsky, M. Ruiz, A. Saxton, B. Schutz, A. Schwing, E. Seidel, S. L. Shapiro, H. Shen, Y. Shen, B. M. Sipőcz, L. Sun, J. Towns, A. Tsokaros, W. Wei, J. Wells, T. J. Williams, J. Xiong, and Z. Zhao. Deep Learning for Multi-Messenger Astrophysics: A Gateway for Discovery in the Big Data Era. *arXiv e-prints*, Feb. 2019.

- [9] I. Arcavi. The First Hours of the GW170817 Kilonova and the Importance of Early Optical and Ultraviolet Observations for Constraining Emission Models. *ApJL*, 855:L23, Mar. 2018.
- [10] S. Ascenzi, M. W. Coughlin, T. Dietrich, R. J. Foley, E. Ramirez-Ruiz, S. Piranomonte, B. Mockler, A. Murguia-Berthier, C. L. Fryer, N. M. Lloyd-Ronning, and S. Rosswog. A luminosity distribution for kilonovae based on short gamma-ray burst afterglows. *ArXiv e-prints*, Nov. 2018.
- [11] A. Bauswein, T. W. Baumgarte, and H.-T. Janka. Prompt Merger Collapse and the Maximum Mass of Neutron Stars. *Physical Review Letters*, 111(13):131101, Sept. 2013.
- [12] A. Bauswein, O. Just, H.-T. Janka, and N. Stergioulas. Neutron-star Radius Constraints from GW170817 and Future Detections. *ApJL*, 850:L34, Dec. 2017.
- [13] P. K. Blanchard, E. Berger, W. Fong, M. Nicholl, J. Leja, C. Conroy, K. D. Alexander, R. Margutti, P. K. G. Williams, Z. Doctor, R. Chornock, V. A. Villar, P. S. Cowperthwaite, J. Annis, D. Brout, D. A. Brown, H.-Y. Chen, T. Eftekhari, J. A. Frieman, D. E. Holz, B. D. Metzger, A. Rest, M. Sako, and M. Soares-Santos. The Electromagnetic Counterpart of the Binary Neutron Star Merger LIGO/Virgo GW170817. VII. Properties of the Host Galaxy and Constraints on the Merger Timescale. *ApJL*, 848:L22, Oct. 2017.
- [14] H.-Y. Chen, M. Fishbach, and D. E. Holz. A two per cent Hubble constant measurement from standard sirens within five years. *Nature*, 562:545–547, Oct. 2018.
- [15] D. F. Chernoff and L. S. Finn. Gravitational radiation, inspiraling binaries, and cosmology. *ApJL*, 411:L5–L8, July 1993.
- [16] M. W. Coughlin, T. Dietrich, B. Margalit, and B. D. Metzger. Multi-messenger Bayesian parameter inference of a binary neutron-star merger. *arXiv e-prints*, page arXiv:1812.04803, Dec 2018.
- [17] D. A. Coulter, R. J. Foley, C. D. Kilpatrick, M. R. Drout, A. L. Piro, B. J. Shappee, M. R. Siebert, J. D. Simon, N. Ulloa, D. Kasen, B. F. Madore, A. Murguia-Berthier, Y.-C. Pan, J. X. Prochaska, E. Ramirez-Ruiz, A. Rest, and C. Rojas-Bravo. Swope Supernova Survey 2017a (SSS17a), the optical counterpart to a gravitational wave source. *Science*, 358:1556–1558, Dec. 2017.
- [18] P. S. Cowperthwaite, E. Berger, V. A. Villar, B. D. Metzger, M. Nicholl, R. Chornock, P. K. Blanchard, W. Fong, R. Margutti, M. Soares-Santos, K. D. Alexander, S. Allam, J. Annis, D. Brout, D. A. Brown, R. E. Butler, H.-Y. Chen, H. T. Diehl, Z. Doctor, M. R. Drout, T. Eftekhari, B. Farr, D. A. Finley, R. J. Foley, J. A. Frieman, C. L. Fryer, J. García-Bellido, M. S. S. Gill, J. Guillochon, K. Herner, D. E. Holz, D. Kasen, R. Kessler, J. Marriner, T. Matheson, E. H. Neilsen, Jr., E. Quataert, A. Palmese, A. Rest, M. Sako, D. M. Scolnic, N. Smith, D. L. Tucker, P. K. G. Williams, E. Balbinot, J. L. Carlin, E. R. Cook, F. Durret, T. S. Li, P. A. A. Lopes, A. C. C. Lourenço, J. L. Marshall, G. E. Medina, J. Muir, R. R. Muñoz, M. Sauseda, D. J. Schlegel, L. F. Secco,

- A. K. Vivas, W. Wester, A. Zenteno, Y. Zhang, T. M. C. Abbott, M. Banerji, K. Bechtol, A. Benoit-Lévy, E. Bertin, E. Buckley-Geer, D. L. Burke, D. Capozzi, A. Carnero Rosell, M. Carrasco Kind, F. J. Castander, M. Croce, C. E. Cunha, C. B. D’Andrea, L. N. da Costa, C. Davis, D. L. DePoy, S. Desai, J. P. Dietrich, A. Drlica-Wagner, T. F. Eifler, A. E. Evrard, E. Fernandez, B. Flaugher, P. Fosalba, E. Gaztanaga, D. W. Gerdes, T. Giannantonio, D. A. Goldstein, D. Gruen, R. A. Gruendl, G. Gutierrez, K. Honscheid, B. Jain, D. J. James, T. Jeltema, M. W. G. Johnson, M. D. Johnson, S. Kent, E. Krause, R. Kron, K. Kuehn, N. Nuropatkin, O. Lahav, M. Lima, H. Lin, M. A. G. Maia, M. March, P. Martini, R. G. McMahon, F. Menanteau, C. J. Miller, R. Miquel, J. J. Mohr, E. Neilsen, R. C. Nichol, R. L. C. Ogando, A. A. Plazas, N. Roe, A. K. Romer, A. Roodman, E. S. Rykoff, E. Sanchez, V. Scarpine, R. Schindler, M. Schubnell, I. Sevilla-Noarbe, M. Smith, R. C. Smith, F. Sobreira, E. Suchyta, M. E. C. Swanson, G. Tarle, D. Thomas, R. C. Thomas, M. A. Troxel, V. Vikram, A. R. Walker, R. H. Wechsler, J. Weller, B. Yanny, and J. Zuntz. The Electromagnetic Counterpart of the Binary Neutron Star Merger LIGO/Virgo GW170817. II. UV, Optical, and Near-infrared Light Curves and Comparison to Kilonova Models. *ApJL*, 848:L17, Oct. 2017.
- [19] P. Creminelli and F. Vernizzi. Dark Energy after GW170817 and GRB170817A. *Physical Review Letters*, 119(25):251302, Dec. 2017.
- [20] N. Dalal, D. E. Holz, S. A. Hughes, and B. Jain. Short GRB and binary black hole standard sirens as a probe of dark energy. , 74(6):063006, Sept. 2006.
- [21] S. De, D. Finstad, J. M. Lattimer, D. A. Brown, E. Berger, and C. M. Biwer. Tidal Deformabilities and Radii of Neutron Stars from the Observation of GW170817. *Physical Review Letters*, 121(9):091102, Aug. 2018.
- [22] M. R. Drout, A. L. Piro, B. J. Shappee, C. D. Kilpatrick, J. D. Simon, C. Contreras, D. A. Coulter, R. J. Foley, M. R. Siebert, N. Morrell, K. Boutsia, F. Di Mille, T. W.-S. Holoién, D. Kasen, J. A. Kollmeier, B. F. Madore, A. J. Monson, A. Murguía-Berthier, Y.-C. Pan, J. X. Prochaska, E. Ramirez-Ruiz, A. Rest, C. Adams, K. Alatalo, E. Bañados, J. Baughman, T. C. Beers, R. A. Bernstein, T. Bitsakis, A. Campillay, T. T. Hansen, C. R. Higgs, A. P. Ji, G. Maravelias, J. L. Marshall, C. Moni Bidin, J. L. Prieto, K. C. Rasmussen, C. Rojas-Bravo, A. L. Strom, N. Ulloa, J. Vargas-González, Z. Wan, and D. D. Whitten. Light curves of the neutron star merger GW170817/SSS17a: Implications for r-process nucleosynthesis. *Science*, 358:1570–1574, Dec. 2017.
- [23] P. A. Evans, S. B. Cenko, J. A. Kennea, S. W. K. Emery, N. P. M. Kuin, O. Korobkin, R. T. Wollaeger, C. L. Fryer, K. K. Madsen, F. A. Harrison, Y. Xu, E. Nakar, K. Hotokezaka, A. Lien, S. Campana, S. R. Oates, E. Troja, A. A. Breeveld, F. E. Marshall, S. D. Barthelmy, A. P. Beardmore, D. N. Burrows, G. Cusumano, A. D’Aì, P. D’Avanzo, V. D’Elia, M. de Pasquale, W. P. Even, C. J. Fontes, K. Forster, J. Garcia, P. Giommi, B. Grefenstette, C. Gronwall, D. H. Hartmann, M. Heida, A. L. Hungerford, M. M. Kasliwal, H. A. Krimm, A. J. Levan, D. Malesani, A. Melandri, H. Miyasaka, J. A. Nousek, P. T. O’Brien, J. P. Osborne, C. Pagani, K. L. Page, D. M. Palmer, M. Perri,

- S. Pike, J. L. Racusin, S. Rosswog, M. H. Siegel, T. Sakamoto, B. Sbarufatti, G. Tagliaferri, N. R. Tanvir, and A. Tohuvavohu. Swift and NuSTAR observations of GW170817: Detection of a blue kilonova. *Science*, 358:1565–1570, Dec. 2017.
- [24] J. M. Ezquiaga and M. Zumalacárregui. Dark Energy After GW170817: Dead Ends and the Road Ahead. *Physical Review Letters*, 119(25):251304, Dec. 2017.
- [25] W. Fong and E. Berger. The Locations of Short Gamma-Ray Bursts as Evidence for Compact Object Binary Progenitors. *ApJ*, 776:18, Oct. 2013.
- [26] W. Fong, E. Berger, R. Chornock, R. Margutti, A. J. Levan, N. R. Tanvir, R. L. Tunnicliffe, I. Czekala, D. B. Fox, D. A. Perley, S. B. Cenko, B. A. Zauderer, T. Laskar, S. E. Persson, A. J. Monson, D. D. Kelson, C. Birk, D. Murphy, M. Servillat, and G. Aniglada. Demographics of the Galaxies Hosting Short-duration Gamma-Ray Bursts. *ApJ*, 769:56, May 2013.
- [27] F. Foucart, M. D. Duez, T. Hinderer, J. Caro, A. R. Williamson, M. Boyle, A. Buonanno, R. Haas, D. A. Hemberger, L. E. Kidder, H. P. Pfeiffer, and M. A. Scheel. Gravitational waveforms from spectral Einstein code simulations: Neutron star-neutron star and low-mass black hole-neutron star binaries. , 99(4):044008, Feb. 2019.
- [28] G. Ghirlanda, O. S. Salafia, Z. Paragi, M. Giroletti, J. Yang, B. Marcote, J. Blanchard, I. Agudo, T. An, M. G. Bernardini, R. Beswick, M. Branchesi, S. Campana, C. Casadio, E. Chassande-Mottin, M. Colpi, S. Covino, P. D’Avanzo, V. D’Elia, S. Frey, M. Gawronski, G. Ghisellini, L. I. Gurvits, P. G. Jonker, H. J. van Langevelde, A. Melandri, J. Moldon, L. Nava, A. Perego, M. A. Perez-Torres, C. Reynolds, R. Salvaterra, G. Tagliaferri, T. Venturi, S. D. Vergani, and M. Zhang. Compact radio emission indicates a structured jet was produced by a binary neutron star merger. *arXiv e-prints*, page arXiv:1808.00469, Aug 2018.
- [29] B. P. Gompertz, A. J. Levan, N. R. Tanvir, J. Hjorth, S. Covino, P. A. Evans, A. S. Fruchter, C. González-Fernández, Z. P. Jin, J. D. Lyman, S. R. Oates, P. T. O’Brien, and K. Wiersema. The Diversity of Kilonova Emission in Short Gamma-Ray Bursts. *ApJ*, 860:62, June 2018.
- [30] G. Hallinan, A. Corsi, K. P. Mooley, K. Hotokezaka, E. Nakar, M. M. Kasliwal, D. L. Kaplan, D. A. Frail, S. T. Myers, T. Murphy, K. De, D. Dobie, J. R. Allison, K. W. Bannister, V. Bhalerao, P. Chandra, T. E. Clarke, S. Giacintucci, A. Y. Q. Ho, A. Horesh, N. E. Kassim, S. R. Kulkarni, E. Lenc, F. J. Lockman, C. Lynch, D. Nichols, S. Nissanke, N. Palliyaguru, W. M. Peters, T. Piran, J. Rana, E. M. Sadler, and L. P. Singer. A radio counterpart to a neutron star merger. *Science*, 358:1579–1583, Dec. 2017.
- [31] J. Hjorth, A. J. Levan, N. R. Tanvir, J. D. Lyman, R. Wojtak, S. L. Schröder, I. Mandel, C. Gall, and S. H. Bruun. The Distance to NGC 4993: The Host Galaxy of the Gravitational-wave Event GW170817. *ApJL*, 848:L31, Oct. 2017.

- [32] K. Hotokezaka, K. Kiuchi, K. Kyutoku, T. Muranushi, Y.-i. Sekiguchi, M. Shibata, and K. Taniguchi. Remnant massive neutron stars of binary neutron star mergers: Evolution process and gravitational waveform. , 88(4):044026, Aug. 2013.
- [33] K. Hotokezaka, E. Nakar, O. Gottlieb, S. Nissanke, K. Masuda, G. Hallinan, K. P. Mooley, and A. T. Deller. A Hubble constant measurement from superluminal motion of the jet in GW170817. *ArXiv e-prints*, June 2018.
- [34] O. Just, A. Bauswein, R. Ardevol Pulpillo, S. Goriely, and H.-T. Janka. Comprehensive nucleosynthesis analysis for ejecta of compact binary mergers. *MNRAS*, 448:541–567, Mar. 2015.
- [35] D. Kasen, N. R. Badnell, and J. Barnes. Opacities and Spectra of the r-process Ejecta from Neutron Star Mergers. *ApJ*, 774:25, Sept. 2013.
- [36] D. Kasen, B. Metzger, J. Barnes, E. Quataert, and E. Ramirez-Ruiz. Origin of the heavy elements in binary neutron-star mergers from a gravitational-wave event. *Nature*, 551:80–84, Nov. 2017.
- [37] M. M. Kasliwal, D. Kasen, R. M. Lau, D. A. Perley, S. Rosswog, E. O. Ofek, K. Hotokezaka, R.-R. Chary, J. Sollerman, A. Goobar, and D. L. Kaplan. Spitzer Mid-Infrared Detections of Neutron Star Merger GW170817 Suggests Synthesis of the Heaviest Elements. *MNRAS*, Jan. 2019.
- [38] M. M. Kasliwal, E. Nakar, L. P. Singer, D. L. Kaplan, D. O. Cook, A. Van Sistine, R. M. Lau, C. Fremling, O. Gottlieb, J. E. Jencson, S. M. Adams, U. Feindt, K. Hotokezaka, S. Ghosh, D. A. Perley, P.-C. Yu, T. Piran, J. R. Allison, G. C. Anupama, A. Balasubramanian, K. W. Bannister, J. Bally, J. Barnes, S. Barway, E. Bellm, V. Bhalerao, D. Bhattacharya, N. Blagorodnova, J. S. Bloom, P. R. Brady, C. Cannella, D. Chatterjee, S. B. Cenko, B. E. Cobb, C. Copperwheat, A. Corsi, K. De, D. Dobie, S. W. K. Emery, P. A. Evans, O. D. Fox, D. A. Frail, C. Frohmaier, A. Goobar, G. Hallinan, F. Harrison, G. Helou, T. Hinderer, A. Y. Q. Ho, A. Horesh, W.-H. Ip, R. Itoh, D. Kasen, H. Kim, N. P. M. Kuin, T. Kupfer, C. Lynch, K. Madsen, P. A. Mazzali, A. A. Miller, K. Mooley, T. Murphy, C.-C. Ngeow, D. Nichols, S. Nissanke, P. Nugent, E. O. Ofek, H. Qi, R. M. Quimby, S. Rosswog, F. Rusu, E. M. Sadler, P. Schmidt, J. Sollerman, I. Steele, A. R. Williamson, Y. Xu, L. Yan, Y. Yatsu, C. Zhang, and W. Zhao. Illuminating gravitational waves: A concordant picture of photons from a neutron star merger. *Science*, 358:1559–1565, Dec. 2017.
- [39] K. Kawaguchi, M. Shibata, and M. Tanaka. Radiative Transfer Simulation for the Optical and Near-infrared Electromagnetic Counterparts to GW170817. *ApJL*, 865:L21, Oct. 2018.
- [40] C. D. Kilpatrick, R. J. Foley, D. Kasen, A. Murguia-Berthier, E. Ramirez-Ruiz, D. A. Coulter, M. R. Drout, A. L. Piro, B. J. Shappee, K. Boutsia, C. Contreras, F. Di Mille, B. F. Madore, N. Morrell, Y.-C. Pan, J. X. Prochaska, A. Rest, C. Rojas-Bravo, M. R. Siebert, J. D. Simon, and N. Ulloa. Electromagnetic Evidence that SSS17a is the Result of a Binary Neutron Star Merger. *Science*, Oct. 2017.

- [41] S. Kisaka, K. Ioka, and E. Nakar. X-Ray-powered Macronovae. *ApJ*, 818:104, Feb. 2016.
- [42] K. Kiuchi, Y. Sekiguchi, M. Shibata, and K. Taniguchi. Long-term general relativistic simulation of binary neutron stars collapsing to a black hole. , 80(6):064037, Sept. 2009.
- [43] K. Kotake. Multiple physical elements to determine the gravitational-wave signatures of core-collapse supernovae. *Comptes Rendus Physique*, 14:318–351, Apr. 2013.
- [44] D. Lazzati, R. Perna, B. J. Morsony, D. Lopez-Camara, M. Cantiello, R. Ciolfi, B. Giacomazzo, and J. C. Workman. Late Time Afterglow Observations Reveal a Collimated Relativistic Jet in the Ejecta of the Binary Neutron Star Merger GW170817. *Physical Review Letters*, 120(24):241103, June 2018.
- [45] A. J. Levan, J. D. Lyman, N. R. Tanvir, J. Hjorth, I. Mandel, E. R. Stanway, D. Steeghs, A. S. Fruchter, E. Troja, S. L. Schröder, K. Wiersema, S. H. Bruun, Z. Cano, S. B. Cenko, A. de Ugarte Postigo, P. Evans, S. Fairhurst, O. D. Fox, J. P. U. Fynbo, B. Gompertz, J. Greiner, M. Im, L. Izzo, P. Jakobsson, T. Kangas, H. G. Khandrika, A. Y. Lien, D. Malesani, P. O’Brien, J. P. Osborne, E. Palazzi, E. Pian, D. A. Perley, S. Rosswog, R. E. Ryan, S. Schulze, P. Sutton, C. C. Thöne, D. J. Watson, and R. A. M. J. Wijers. The Environment of the Binary Neutron Star Merger GW170817. *ApJL*, 848:L28, Oct. 2017.
- [46] J. Logue, C. D. Ott, I. S. Heng, P. Kalmus, and J. H. C. Scargill. Inferring core-collapse supernova physics with gravitational waves. , 86:044023, Aug 2012.
- [47] I. Mandel, W. M. Farr, and J. R. Gair. Extracting distribution parameters from multiple uncertain observations with selection biases. *ArXiv e-prints*, Sept. 2018.
- [48] B. Margalit and B. D. Metzger. Constraining the Maximum Mass of Neutron Stars from Multi-messenger Observations of GW170817. *ApJL*, 850:L19, Dec. 2017.
- [49] B. D. Metzger and R. Fernández. Red or blue? A potential kilonova imprint of the delay until black hole formation following a neutron star merger. *MNRAS*, 441:3444–3453, July 2014.
- [50] B. D. Metzger, T. A. Thompson, and E. Quataert. A Magnetar Origin for the Kilonova Ejecta in GW170817. *ApJ*, 856:101, Apr. 2018.
- [51] K. P. Mooley, A. T. Deller, O. Gottlieb, E. Nakar, G. Hallinan, S. Bourke, D. A. Frail, A. Horesh, A. Corsi, and K. Hotokezaka. Superluminal motion of a relativistic jet in the neutron-star merger GW170817. *Nature*, 561:355–359, Sept. 2018.
- [52] K. P. Mooley, D. A. Frail, D. Dobie, E. Lenc, A. Corsi, K. De, A. J. Nayana, S. Makhathini, I. Heywood, T. Murphy, D. L. Kaplan, P. Chandra, O. Smirnov, E. Nakar, G. Hallinan, F. Camilo, R. Fender, S. Goedhart, P. Groot, M. M. Kasliwal, S. R. Kulkarni, and P. A. Woudt. A Strong Jet Signature in the Late-time Light Curve of GW170817. *ApJL*, 868:L11, Nov. 2018.

- [53] S. Nissanke, D. E. Holz, N. Dalal, S. A. Hughes, J. L. Sievers, and C. M. Hirata. Determining the Hubble constant from gravitational wave observations of merging compact binaries. *arXiv e-prints*, July 2013.
- [54] A. Palmese, W. Hartley, F. Tarsitano, C. Conselice, O. Lahav, S. Allam, J. Annis, H. Lin, M. Soares-Santos, D. Tucker, D. Brout, M. Banerji, K. Bechtol, H. T. Diehl, A. Fruchter, J. García-Bellido, K. Herner, A. J. Levan, T. S. Li, C. Lidman, K. Misra, M. Sako, D. Scolnic, M. Smith, T. M. C. Abbott, F. B. Abdalla, A. Benoit-Lévy, E. Bertin, D. Brooks, E. Buckley-Geer, A. Carnero Rosell, M. Carrasco Kind, J. Carretero, F. J. Castander, C. E. Cunha, C. B. D’Andrea, L. N. da Costa, C. Davis, D. L. DePoy, S. Desai, J. P. Dietrich, P. Doel, A. Drlica-Wagner, T. F. Eifler, A. E. Evrard, B. Flaugher, P. Fosalba, J. Frieman, E. Gaztanaga, D. W. Gerdes, T. Giannantonio, D. Gruen, R. A. Gruendl, J. Gschwend, G. Gutierrez, K. Honscheid, B. Jain, D. J. James, T. Jeltema, M. W. G. Johnson, M. D. Johnson, E. Krause, R. Kron, K. Kuehn, S. Kuhlmann, N. Kuropatkin, M. Lima, M. A. G. Maia, M. March, J. L. Marshall, R. G. McMahon, F. Menanteau, C. J. Miller, R. Miquel, E. Neilsen, R. L. C. Ogando, A. A. Plazas, K. Reil, A. K. Romer, E. Sanchez, R. Schindler, R. C. Smith, F. Sobreira, E. Suchyta, M. E. C. Swanson, G. Tarle, D. Thomas, R. C. Thomas, A. R. Walker, J. Weller, Y. Zhang, and J. Zuntz. Evidence for Dynamically Driven Formation of the GW170817 Neutron Star Binary in NGC 4993. *ApJL*, 849:L34, Nov. 2017.
- [55] Y.-C. Pan, C. D. Kilpatrick, J. D. Simon, E. Xhakaj, K. Boutsia, D. A. Coulter, M. R. Drout, R. J. Foley, D. Kasen, N. Morrell, A. Murguia-Berthier, D. Osip, A. L. Piro, J. X. Prochaska, E. Ramirez-Ruiz, A. Rest, C. Rojas-Bravo, B. J. Shappee, and M. R. Siebert. The Old Host-galaxy Environment of SSS17a, the First Electromagnetic Counterpart to a Gravitational-wave Source. *ApJL*, 848:L30, Oct. 2017.
- [56] K. Pardo, M. Fishbach, D. E. Holz, and D. N. Spergel. Limits on the number of spacetime dimensions from GW170817. , 7:048, July 2018.
- [57] V. Paschalidis, M. Ruiz, and S. L. Shapiro. Relativistic Simulations of Black Hole-Neutron Star Coalescence: The Jet Emerges. *ApJL*, 806:L14, June 2015.
- [58] E. Pian, P. D’Avanzo, S. Benetti, M. Branchesi, E. Brocato, S. Campana, E. Cappellaro, S. Covino, V. D’Elia, J. P. U. Fynbo, F. Getman, G. Ghirlanda, G. Ghisellini, A. Grado, G. Greco, J. Hjorth, C. Kouveliotou, A. Levan, L. Limatola, D. Malesani, P. A. Mazzali, A. Melandri, P. Møller, L. Nicastro, E. Palazzi, S. Piranomonte, A. Rossi, O. S. Salafia, J. Selsing, G. Stratta, M. Tanaka, N. R. Tanvir, L. Tomasella, D. Watson, S. Yang, L. Amati, L. A. Antonelli, S. Ascenzi, M. G. Bernardini, M. Boër, F. Bufano, A. Bulgarelli, M. Capaccioli, P. Casella, A. J. Castro-Tirado, E. Chassande-Mottin, R. Ciolfi, C. M. Copperwheat, M. Dadina, G. De Cesare, A. di Paola, Y. Z. Fan, B. Gendre, G. Giuffrida, A. Giunta, L. K. Hunt, G. L. Israel, Z.-P. Jin, M. M. Kasliwal, S. Klose, M. Lisi, F. Longo, E. Maiorano, M. Mapelli, N. Masetti, L. Nava, B. Patricelli, D. Perley, A. Pescalli, T. Piran, A. Possenti, L. Pulone, M. Razzano, R. Salvaterra, P. Schipani, M. Spera, A. Stamerra, L. Stella, G. Tagliaferri, V. Testa, E. Troja, M. Turatto, S. D. Vergani, and D. Vergani. Spectroscopic identification of r-process nucleosynthesis in a double neutron-star merger. *Nature*, 551:67–70, Nov. 2017.

- [59] A. L. Piro and J. A. Kollmeier. Evidence for Cocoon Emission from the Early Light Curve of SSS17a. *ApJ*, 855:103, Mar. 2018.
- [60] Planck Collaboration, N. Aghanim, Y. Akrami, M. Ashdown, J. Aumont, C. Baccigalupi, M. Ballardini, A. J. Banday, R. B. Barreiro, N. Bartolo, S. Basak, R. Battye, K. Benabed, J.-P. Bernard, M. Bersanelli, P. Bielewicz, J. J. Bock, J. R. Bond, J. Borrill, F. R. Bouchet, F. Boulanger, M. Bucher, C. Burigana, R. C. Butler, E. Calabrese, J.-F. Cardoso, J. Carron, A. Challinor, H. C. Chiang, J. Chluba, L. P. L. Colombo, C. Combet, D. Contreras, B. P. Crill, F. Cuttaia, P. de Bernardis, G. de Zotti, J. Delabrouille, J.-M. Delouis, E. Di Valentino, J. M. Diego, O. Doré, M. Douspis, A. Ducout, X. Dupac, S. Dusini, G. Efstathiou, F. Elsner, T. A. Enßlin, H. K. Eriksen, Y. Fantaye, M. Farhang, J. Fergusson, R. Fernandez-Cobos, F. Finelli, F. Forastieri, M. Frailis, E. Franceschi, A. Frolov, S. Galeotta, S. Galli, K. Ganga, R. T. Génova-Santos, M. Gerbino, T. Ghosh, J. González-Nuevo, K. M. Górski, S. Gratton, A. Gruppuso, J. E. Gudmundsson, J. Hamann, W. Handley, D. Herranz, E. Hivon, Z. Huang, A. H. Jaffe, W. C. Jones, A. Karakci, E. Keihänen, R. Kesitalo, K. Kiiveri, J. Kim, T. S. Kisner, L. Knox, N. Krachmalnicoff, M. Kunz, H. Kurki-Suonio, G. Lagache, J.-M. Lamarre, A. Lasenby, M. Lattanzi, C. R. Lawrence, M. Le Jeune, P. Lemos, J. Lesgourgues, F. Levrier, A. Lewis, M. Liguori, P. B. Lilje, M. Lilley, V. Lindholm, M. López-Caniego, P. M. Lubin, Y.-Z. Ma, J. F. Macías-Pérez, G. Maggio, D. Maino, N. Mandolesi, A. Mangilli, A. Marcos-Caballero, M. Maris, P. G. Martin, M. Martinelli, E. Martínez-González, S. Matarrese, N. Mauri, J. D. McEwen, P. R. Meinhold, A. Melchiorri, A. Mennella, M. Migliaccio, M. Millea, S. Mitra, M.-A. Miville-Deschênes, D. Molinari, L. Montier, G. Morgante, A. Moss, P. Natoli, H. U. Nørgaard-Nielsen, L. Pagano, D. Paoletti, B. Partridge, G. Patanchon, H. V. Peiris, F. Perrotta, V. Pettorino, F. Piacentini, L. Polastri, G. Polenta, J.-L. Puget, J. P. Rachen, M. Reinecke, M. Remazeilles, A. Renzi, G. Rocha, C. Rosset, G. Roudier, J. A. Rubiño-Martín, B. Ruiz-Granados, L. Salvati, M. Sandri, M. Savelainen, D. Scott, E. P. S. Shellard, C. Sirignano, G. Sirri, L. D. Spencer, R. Sunyaev, A.-S. Suur-Uski, J. A. Tauber, D. Tavagnacco, M. Tenti, L. Toffolatti, M. Tomasi, T. Trombetti, L. Valenziano, J. Valiviita, B. Van Tent, L. Vibert, P. Vielva, F. Villa, N. Vittorio, B. D. Wandelt, I. K. Wehus, M. White, S. D. M. White, A. Zacchei, and A. Zonca. Planck 2018 results. VI. Cosmological parameters. *ArXiv e-prints*, July 2018.
- [61] J. Powell and B. Müller. Gravitational Wave Emission from 3D Explosion Models of Core-Collapse Supernovae with Low and Normal Explosion Energies. *arXiv e-prints*, page arXiv:1812.05738, Dec 2018.
- [62] J. X. Prochaska, J. S. Bloom, H.-W. Chen, R. J. Foley, D. A. Perley, E. Ramirez-Ruiz, J. Granot, W. H. Lee, D. Pooley, K. Alatalo, K. Hurley, M. C. Cooper, A. K. Dupree, B. F. Gerke, B. M. S. Hansen, J. S. Kalirai, J. A. Newman, R. M. Rich, H. Richer, S. A. Stanford, D. Stern, and W. J. M. van Breugel. The Galaxy Hosts and Large-Scale Environments of Short-Hard Gamma-Ray Bursts. *ApJ*, 642:989–994, May 2006.
- [63] D. Radice and L. Dai. Multimessenger Parameter Estimation of GW170817. *arXiv e-prints*, page arXiv:1810.12917, Oct 2018.

- [64] A. G. Riess, S. Casertano, W. Yuan, L. Macri, B. Bucciarelli, M. G. Lattanzi, J. W. MacKenty, J. B. Bowers, W. Zheng, A. V. Filippenko, C. Huang, and R. I. Anderson. Milky Way Cepheid Standards for Measuring Cosmic Distances and Application to Gaia DR2: Implications for the Hubble Constant. *ApJ*, 861:126, July 2018.
- [65] S. Rosswog and M. B. Davies. High-resolution calculations of merging neutron stars - I. Model description and hydrodynamic evolution. *MNRAS*, 334:481–497, Aug. 2002.
- [66] S. Rosswog, M. Liebendörfer, F.-K. Thielemann, M. B. Davies, W. Benz, and T. Piran. Mass ejection in neutron star mergers. *A&A*, 341:499–526, Jan. 1999.
- [67] S. Rosswog, J. Sollerman, U. Feindt, A. Goobar, O. Korobkin, R. Wollaeger, C. Fremling, and M. M. Kasliwal. The first direct double neutron star merger detection: Implications for cosmic nucleosynthesis. *A&A*, 615:A132, July 2018.
- [68] M. Ruiz, S. L. Shapiro, and A. Tsokaros. GW170817, general relativistic magnetohydrodynamic simulations, and the neutron star maximum mass. *Phys. Rev. Lett.*, 97(2):021501, Jan. 2018.
- [69] K. Scholberg. Supernova Neutrino Detection. *Annual Review of Nuclear and Particle Science*, 62:81–103, Nov. 2012.
- [70] B. F. Schutz. Determining the Hubble constant from gravitational wave observations. *Nature*, 323:310, Sept. 1986.
- [71] M. Shibata, S. Fujibayashi, K. Hotokezaka, K. Kiuchi, K. Kyutoku, Y. Sekiguchi, and M. Tanaka. Modeling GW170817 based on numerical relativity and its implications. *Phys. Rev. Lett.*, 96(12):123012, Dec. 2017.
- [72] S. J. Smartt, T.-W. Chen, A. Jerkstrand, M. Coughlin, E. Kankare, S. A. Sim, M. Fraser, C. Inerra, K. Maguire, K. C. Chambers, M. E. Huber, T. Krühler, G. Leloudas, M. Magee, L. J. Shingles, K. W. Smith, D. R. Young, J. Tonry, R. Kotak, A. Gal-Yam, J. D. Lyman, D. S. Homan, C. Agliozzo, J. P. Anderson, C. R. Angus, C. Ashall, C. Barbarino, F. E. Bauer, M. Berton, M. T. Botticella, M. Bulla, J. Bulger, G. Cannizzaro, Z. Cano, R. Cartier, A. Cikota, P. Clark, A. De Cia, M. Della Valle, L. Denneau, M. Dennefeld, L. Dessart, G. Dimitriadis, N. Elias-Rosa, R. E. Firth, H. Flewelling, A. Flörs, A. Franckowiak, C. Frohmaier, L. Galbany, S. González-Gaitán, J. Greiner, M. Gromadzki, A. N. Guelbenzu, C. P. Gutiérrez, A. Hamanowicz, L. Hanlon, J. Harnanen, K. E. Heintz, A. Heinze, M.-S. Hernandez, S. T. Hodgkin, I. M. Hook, L. Izzo, P. A. James, P. G. Jonker, W. E. Kerzendorf, S. Klose, Z. Kostrzewa-Rutkowska, M. Kowalski, M. Kromer, H. Kuncarayakti, A. Lawrence, T. B. Lowe, E. A. Magnier, I. Manulis, A. Martin-Carrillo, S. Mattila, O. McBrien, A. Müller, J. Nordin, D. O’Neill, F. Onori, J. T. Palmerio, A. Pastorello, F. Patat, G. Pignata, P. Podsiadlowski, M. L. Pumo, S. J. Prentice, A. Rau, A. Razza, A. Rest, T. Reynolds, R. Roy, A. J. Ruiter, K. A. Rybicki, L. Salmon, P. Schady, A. S. B. Schultz, T. Schweyer, I. R. Seitenzahl, M. Smith, J. Sollerman, B. Stalder, C. W. Stubbs, M. Sullivan, H. Szegedi, F. Taddia, S. Taubenberger, G. Terreran, B. van Soelen, J. Vos, R. J. Wainscoat, N. A. Walton, C. Waters,

- H. Weiland, M. Willman, P. Wiseman, D. E. Wright, L. Wyrzykowski, and O. Yaron. A kilonova as the electromagnetic counterpart to a gravitational-wave source. *Nature*, 551:75–79, Nov. 2017.
- [73] N. R. Tanvir, A. J. Levan, C. González-Fernández, O. Korobkin, I. Mandel, S. Rosswog, J. Hjorth, P. D’Avanzo, A. S. Fruchter, C. L. Fryer, T. Kangas, B. Milvang-Jensen, S. Rosetti, D. Steeghs, R. T. Wollaeger, Z. Cano, C. M. Copperwheat, S. Covino, V. D’Elia, A. de Ugarte Postigo, P. A. Evans, W. P. Even, S. Fairhurst, R. Figuera Jaimes, C. J. Fontes, Y. I. Fujii, J. P. U. Fynbo, B. P. Gompertz, J. Greiner, G. Hodosan, M. J. Irwin, P. Jakobsson, U. G. Jørgensen, D. A. Kann, J. D. Lyman, D. Malesani, R. G. McMahon, A. Melandri, P. T. O’Brien, J. P. Osborne, E. Palazzi, D. A. Perley, E. Pian, S. Piranomonte, M. Rabus, E. Rol, A. Rowlinson, S. Schulze, P. Sutton, C. C. Thöne, K. Ulaczyk, D. Watson, K. Wiersema, and R. A. M. J. Wijers. The Emergence of a Lanthanide-rich Kilonova Following the Merger of Two Neutron Stars. *ApJL*, 848:L27, Oct. 2017.
- [74] The LIGO Scientific Collaboration, the Virgo Collaboration, B. P. Abbott, R. Abbott, T. D. Abbott, S. Abraham, F. Acernese, K. Ackley, C. Adams, R. X. Adhikari, and et al. GWTC-1: A Gravitational-Wave Transient Catalog of Compact Binary Mergers Observed by LIGO and Virgo during the First and Second Observing Runs. *arXiv e-prints*, Nov. 2018.
- [75] E. Troja, L. Piro, H. van Eerten, R. T. Wollaeger, M. Im, O. D. Fox, N. R. Butler, S. B. Cenko, T. Sakamoto, C. L. Fryer, R. Ricci, A. Lien, R. E. Ryan, O. Korobkin, S.-K. Lee, J. M. Burgess, W. H. Lee, A. M. Watson, C. Choi, S. Covino, P. D’Avanzo, C. J. Fontes, J. B. González, H. G. Khandrika, J. Kim, S.-L. Kim, C.-U. Lee, H. M. Lee, A. Kutyrev, G. Lim, R. Sánchez-Ramírez, S. Veilleux, M. H. Wieringa, and Y. Yoon. The X-ray counterpart to the gravitational-wave event GW170817. *Nature*, 551:71–74, Nov. 2017.
- [76] V. A. Villar, J. Guillochon, E. Berger, B. D. Metzger, P. S. Cowperthwaite, M. Nicholl, K. D. Alexander, P. K. Blanchard, R. Chornock, T. Eftekhari, W. Fong, R. Margutti, and P. K. G. Williams. The Combined Ultraviolet, Optical, and Near-infrared Light Curves of the Kilonova Associated with the Binary Neutron Star Merger GW170817: Unified Data Set, Analytic Models, and Physical Implications. *ApJL*, 851:L21, Dec. 2017.
- [77] E. Waxman, E. O. Ofek, D. Kushnir, and A. Gal-Yam. Constraints on the ejecta of the GW170817 neutron star merger from its electromagnetic emission. *MNRAS*, 481:3423–3441, Dec. 2018.
- [78] M.-R. Wu, J. Barnes, G. Martinez-Pinedo, and B. D. Metzger. Fingerprints of heavy element nucleosynthesis in the late-time lightcurves of kilonovae. *ArXiv e-prints*, Aug. 2018.
- [79] Y.-W. Yu, L.-D. Liu, and Z.-G. Dai. A Long-lived Remnant Neutron Star after GW170817 Inferred from Its Associated Kilonova. *ApJ*, 861:114, July 2018.

- [80] Y. Zhu, R. T. Wollaeger, N. Vassh, R. Surman, T. M. Sprouse, M. R. Mumpower, P. Möller, G. C. McLaughlin, O. Korobkin, T. Kawano, P. J. Jaffke, E. M. Holmbeck, C. L. Fryer, W. P. Even, A. J. Couture, and J. Barnes. Californium-254 and Kilonova Light Curves. *ApJL*, 863:L23, Aug. 2018.